# A Multi-Phase Transport study of $J/\psi$ suppression at RHIC

Bin Zhang [a,b] C.M. Ko [b] Bao-An Li [a] Ziwei Lin [b] Ben-Hao Sa [b]

[a] *Arkansas State University*

[b] *Texas A&M University*

*Presented by:* Bin Zhang

**Abstract**

$J/\psi$ suppression is one of the most promising signals of Quark-Gluon-Plasma formation. In an infinite Quark-Gluon-Plasma, due to color screening, $J/\psi$'s are dissociated into free $c\bar{c}$ pairs. As Quark-Gluon-Plasma produced in relativistic nucleus-nucleus collisions have finite size and finite life time, a study of $J/\psi$ suppression in realistic space-time picture is necessary. We performed such a study using A Multi-Phase Transport model that includes both parton and hadron space-time evolution. Besides plasma dissociation, we included dissociation of $J/\psi$ by collisions with gluons and hadrons. Both plasma screening and gluon scatterings are found to be important for $J/\psi$ suppression in Au+Au collisions, and the effect due to absorption by hadrons is minor. The final $J/\psi$ survival probability is only a few percent. For S+S collisions, in spite of the initial high parton density, due to short plasma life time, the plasma screening dissociation is absent. This provides an opportunity for the study of gluon-$J/\psi$ interactions.

*Preprint submitted to Elsevier Preprint* *20 December 2000*